

WORRY FREE® by Lilly Miller®
Ferramol™
Slug&Snail BAIT

WHY WORRY FREE® SLUG AND SNAIL BAIT IS EFFECTIVE

This product is a unique blend of an iron phosphate active ingredient, originating from soil, with slug and snail bait additives. It is used as an ingredient in fertilizers. The bait which is not ingested by snails and slugs will degrade and become a part of the soil.

The bait is attractive to slugs and snails and lures them from their hiding places and plants. Ingestion, even in small amounts, will cause them to cease feeding. This physiological effect of the bait gives immediate protection to the plants even though the slugs and snails may remain in the area. After eating the bait, the slugs and snails cease feeding, become less mobile and begin to die within three to six days. Dead slugs and snails may not be visible as they often crawl away to secluded places to die. Plant protection will be observed in the decrease in plant damage.

This product is effective against a wide variety of slugs and snails and will give protection to home lawns, gardens, greenhouses, outdoor ornamentals, vegetable gardens, fruits, berries, citrus and crop plants. The bait can be scattered on the lawn or on the soil around any vegetable plants, flowers or fruit trees or bushes to be protected.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For household home garden use.

KILLS SLUGS & SNAILS

HOW TO APPLY: The slug and snail bait granules should be scattered on the soil around or near the plants to be protected. For broadcast application standard broadcast spreaders may be used. For row application standard granular spreaders may be used. Apply bait evenly at approximately 1 lb. per 1000 square feet (0.15 oz., or about 1 level teaspoon, per square yard) and reapply as the bait is consumed or at least every two weeks. Do not place in piles. If the ground is dry, wet it before applying bait. The soil should be moist but with little or no standing water.

Reapply as the bait is consumed or at least every two weeks. Apply more heavily if the infestation is severe, if the area is heavily watered or after long periods of heavy rain. See specific directions for different plant types and for inside greenhouses.

WHEN TO APPLY: Evening is the best time to apply the bait, as slugs and snails travel and feed mostly by night or early morning.

WHERE TO APPLY: All likely areas of infestation should be treated, especially around the perimeter of garden plots because these pests travel into plant areas from daytime refuges. They favor damp places around vegetable plants such as beans, tomatoes, lettuce, cabbage, celery and squash. Other favorite areas are flower gardens, rockeries, hedges, dichondra lawns, citrus groves, ivy patches, and other ground cover where they obtain shelter by day.

VEGETABLES: The bait can be used to protect any vegetables from slug and snail damage. Scatter the bait around the perimeter of the vegetable plot at approximately 1 lb. per 1000 square feet to provide a protective "barrier" for slugs and snails entering the garden plot. If slugs or snails are inside the rows, then scatter the bait on the soil around the base of the plants and between the rows.

FRUITS INCLUDING CITRUS: The bait can be used to protect fruits from slugs and snails. For seedlings spread the bait around the base of the stem. Apply at 0.15 oz., or 1 level teaspoon, per square yard, in a 6 inch circular band around the base of the plants to be protected. For older trees, spread the bait around the base of the tree to intercept slugs and snails traveling to the trunk. Apply the bait at approximately 1 lb. per 1000 square feet for orchards using standard fertilizer granular spreaders.

BERRIES: The bait can be used to protect berries from slugs and snails. Spread the bait around the perimeter of the plot to intercept slugs and snails migrating toward the berries. Use a rate of approximately 1 lb. per 1000 square feet and scatter by hand or with granular spreaders. If slugs and snails are already in the plots, then carefully spread bait between the furrows near the base of the plants. For small plots, treat around the base of the plants to be protected. Do not spread over the entire area but apply selectively.

GREENHOUSES: Where slugs or snails are a problem in greenhouses, scatter the bait in the plant pots of plants being damaged or around pots on greenhouse benches. Apply about 1/2 teaspoon per 9 inch pot.

OUTDOOR ORNAMENTALS: Scatter bait in a 6 inch circular band around the base of the plants to be protected at 0.15 oz., or 1 level teaspoon, per square yard. If plants are next to a grassy area, spread the bait between the ornamentals and the grass. Slugs and snails traveling to the plants will encounter the bait before reaching the plant. In these situations, scatter the bait around the perimeter of the plot at approximately 1 lb. per 1000 square feet to intercept snails and slugs traveling to the plot.

LAWNS: The bait can be used to protect lawns. When slugs or snails are detected, scatter the bait at a rate of approximately 1 lb. per 1000 square feet. Scatter by hand or with a granular spreader where the slugs or snails are observed.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. PESTICIDE STORAGE: Store this product in its original container and keep in a secure storage area out of reach of children and domestic animals. CONTAINER DISPOSAL: **If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID: If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS: For terrestrial uses. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

OUR COMMITMENT TO YOUR SATISFACTION:
If you are not happy with the way this product works in your garden, please let us know. We will be happy to replace the product or refund your purchase price. Simply send your name, address and proof of purchase to **LILLY MILLER BRANDS,** PO Box 2289, Clackamas, Oregon 97015.

0 70624 10818 5
08801081 - 09/04

Distributed by:
LILLY MILLER BRANDS
PO Box 2289
Clackamas, Oregon 97015

EPA Reg. No. 67702-3-33116
EPA Est. 35497-OR-1

This package is sold by weight.
Contents may have settled during shipment.

www.lillymiller.com

Manufactured by W. Neudorff GmbH KG

NEUDORFF®

FERRAMOL™ is a trademark of W. Neudorff GmbH KG
U.S. Patent # 5,437,870
Made in Germany.